## Trends in Parasitology | Parasite of the Month

# Dermanyssus gallinae (Poultry red mite)

David Hartmann (D<sup>1</sup>, Jan Votýpka (D<sup>1,2</sup>, Julius Lukeš (D<sup>1,3</sup>, and Jan Perner (D<sup>1,\*</sup>

<sup>1</sup>Institute of Parasitology, Biology Centre, Czech Academy of Sciences, České Budějovice (Budweis), Czech Republic
<sup>2</sup>Department of Parasitology, Faculty of Science, Charles University, Prague, Czech Republic
<sup>3</sup>Faculty of Science, University of South Bohemia, České Budějovice (Budweis), Czech Republic



The poultry red mite, *Dermanyssus gallinae*, is a global pest of free-living and farmed birds. As a blood-feeding ectoparasite, it particularly affects the welfare of farmed egg-laying hens. Under optimal conditions, especially in the summer, the life cycle of the parasites can be completed within a week, resulting in their population surges. Heavily infested farms display decreased egg production and economic losses for the poultry industry. The ability of mites to survive for extended periods outside of the host is a major challenge in controlling efforts in poultry houses. Integrated Pest Management strategies are deployed as countermeasures to curb the population of mites. Besides farmed poultry, *D. gallinae* is also a significant nuisance to pet birds, especially parrots and pigeons. In addition to its impact on avian health, this mite is a public health concern due to its role as a vector of pathogens, and its medical impact (gamasoidosis, also known as dermanyssosis) on humans.



### **KEY FACTS:**

The poultry red mite is the most troublesome parasite of egg-laying hens globally, excluding North America, Indonesia, and New Zealand, where the poultry is predominantly parasitised by the northern fowl mite (*Ornithonyssus sylviarum*).

It has a direct life cycle with non-feeding larvae, and blood-feeding protonymphs, deutonymphs, and adults. Bloodfeeding, that can take up to 60 min, negatively affects bird health, welfare, and egg production, resulting in significant economic losses to the poultry industry.

Each female can lay eggs multiple times after each blood-feeding, contributing to rapid population growth.

Hens are primarily attacked at night, when they can be victims to thousands of mites at various stages of development.

### DISEASE FACTS:

Feeding of *D. gallinae* mites can cause dermatitis and anemia in infested birds, leading to significant discomfort, and overall poor health.

Rarely, *D. gallinae* can also feed on humans, causing mild to severe dermatitis.

The stable midgut bacterial microbiome of *D. gallinae*, which includes genera such as *Bartonella*, *Staphylococcus*, *Rickettsiella*, and Micrococcaceae, may contribute to the mite's role as a significant vector and reservoir for pathogens affecting poultry and posing zoonotic risks to human health.

#### TAXONOMY AND CLASSIFICATION:

PHYLUM: Arthropoda CLASS: Arachnida ORDER: Mesostigmata FAMILY: Dermanyssidae GENUS: Dermanyssus SPECIES: D. gallinae (de Geer, 1778)

\*Correspondence: perner@paru.cas.cz (J. Perner).

https://doi.org/10.1016/j.pt.2024.08.006 1055



Trends in Parasitology, November 2024, Vol. 40, No. 11

© 2024 Elsevier Ltd. All rights are reserved, including those for text and data mining, Al training, and similar technologies.

## Trends in Parasitology | Parasite of the Month

### Acknowledgments

This work was supported by the Czech Science Foundation (No: 22-18424M) and the Technology Agency of the Czech Republic (TREND Programme FW10010308).

### **Declaration of interests**

The authors declare no competing interests.

### Resources

www.fao.org/faostat/en/#data/QCL

### Literature

- 1. Banović, P. et al. (2024) One health approach to study human health risks associated with Dermanyssus gallinae mites. Heliyon 10, e30539
- 2. Decru, E. et al. (2020) Possibilities for IPM strategies in European laying hen farms for improved control of the poultry red mite (*Dermanyssus gallinae*): details and state of affairs. Front. Vet. Sci. 7, 565866
- 3. George, D.R. et al. (2015) Should the poultry red mite Dermanyssus gallinae be of wider concern for veterinary and medical science? Parasit. Vectors 8, 178
- 4. Kavallari, A. et al. (2018) Avian mite dermatitis: diagnostic challenges and unmet needs. Parasite Immunol. 40, e12539
- 5. Ribeiro, J. et al. (2023) Blood-feeding adaptations and virome assessment of the poultry red mite Dermanyssus gallinae guided by RNA-seq. Commun. Biol. 6, 517
- 6. Schiavone, A. et al. (2022) Dermanyssus gallinae: the long journey of the poultry red mite to become a vector. Parasit. Vectors 15, 29
- 7. Silva, G.G.d. et al. (2023) Advances in non-chemical tools to control poultry hematophagous mites. Vet. Sci. 10, 589
- 8. Sparagano, O. *et al.* (2020) *Dermanyssus gallinae* and chicken egg production: impact, management, and a predicted compatibility matrix for integrated approaches. *Exp. Appl. Acarol.* 82, 441–453
- 9. Sparagano, O. and Ho, J. (2020) Parasitic mite fauna in Asian poultry farming systems. Front. Vet. Sci. 7, 400
- 10. Wu-Chuang, A. et al. (2023) Variation of bacterial community assembly over developmental stages and midgut of Dermanyssus gallinae. Microb. Ecol. 86, 2400–2413

